



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07K 16/00, 17/00, C07H 21/02, 21/04, A61K 39/395	A1	(11) International Publication Number: WO 98/50432 (43) International Publication Date: 12 November 1998 (12.11.98)
(21) International Application Number: PCT/US98/08840 (22) International Filing Date: 30 April 1998 (30.04.98) (30) Priority Data: 60/045,888 7 May 1997 (07.05.97) . US 09/070,637 30 April 1998 (30.04.98) US (71) Applicant (for all designated States except US): BRIS- TOL-MYERS SQUIBB COMPANY [US/US]; 345 Park Avenue, New York, NY 10154 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): SIEMERS, Nathan, O. [US/US]; 120 Austin Avenue, Hamilton Township, NJ 08629 (US). YARNOLD, Susan [US/US]; 2342 47th Av- enue S.W., Seattle, WA 98116 (US). SENTER, Peter, D. [US/US]; 9000 40th Avenue N.E., Seattle, WA 98116 (US). (74) Agents: PARMELEE, Steven, W. et al.; Townsend and Townsend and Crew LLP, 8th floor, Two Embarcadero Center, San Francisco, CA 94111 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: RECOMBINANT ANTIBODY-ENZYME FUSION PROTEINS (57) Abstract The invention relates to recombinantly produced fusion polypeptides comprising antibody V _H and V _L sequences operatively linked to a β -lactamase for use in the delivery of cytotoxic drugs to tumor cells.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

RECOMBINANT ANTIBODY-ENZYME FUSION PROTEINS

5

BACKGROUND OF THE INVENTION

A considerable amount of attention has been directed towards the use of monoclonal antibody-enzyme conjugates in combination with suitable prodrugs for the selective delivery of chemotherapeutic agents to tumors (reviewed in Senter et al., Bioconjugate Chem., 4:3-9 (1993); Jungheim et al., Chem Rev., 94:1553-1566 (1994); Bagshawe et al., Ann. Oncol., 5:879-891 (1994)). The monoclonal antibody (mAb) portions of these immunoconjugates recognize tumor-selective antigens and are capable of delivering the enzymes to tumor masses. Once tumor localization and systemic conjugate clearance has taken place, a non-cytotoxic prodrug form of a chemotherapeutic drug is administered which is converted into an active drug by the targeted enzyme. This leads to the selective delivery of anticancer drugs to sites of neoplasia. Pharmacokinetic studies have shown that the intratumoral drug concentrations resulting from mAb-enzyme/prodrug combinations can be significantly greater than that achieved by systemic drug administration (Bosslet et al., Cancer Res., 54:2151-2159 (1994); Svensson et al., Cancer Res., 55:2357-2365 (1995); Wallace et al., Cancer Res., 54:2719-2327 (1994)). This probably accounts for the observed antitumor activities, which include complete tumor regressions and cures in a number of different models for human cancer (Springer et al., Eur. J. Cancer, 27:1361-1366 (1991); Meyer et al., Cancer Res., 53:3956-3963 (1993); Eccles et al., Cancer Res., 54:5171-5177 (1994); Kerr et al., Cancer Res., 55:3558-3563 (1995)).

Kerr et al. disclosed the use of antibodies against the human p97 (melanotransferrin) tumor antigen for the delivery of β -lactamase (bL) to tumor cells (Kerr et al., Cancer Res., 55:3558-3563 (1995)). This antigen has been found to be overexpressed on a majority of clinical melanoma isolates and is also observed on many human carcinomas

(Woodbury et al., Proc. Natl. Acad. Sci. (USA), 77:2183-2187 (1980); Brown et al., J. Immun., 127:539-545 (1981), Brown et al., Proc. Natl. Acad. Sci. (USA), 78:539-543 (1981); Rose et al., Proc. Natl. Acad. Sci. (USA) (1986)). Significant
5 antitumor activities have been obtained using the combination of a chemically produced anti p97-Fab'-bL conjugate in combination with CCM (7-(4-carboxybutan-amido)cephalosporin mustard; Kerr et al., Cancer Res., 55:3558-3563 (1995)), a
10 cephalosporin containing prodrug of phenylenediamine mustard (PDM). These effects were observed in a melanoma tumor model that was resistant to the activities of PDM.

A major concern in the development of mAb-bL conjugates for clinical testing is conjugate uniformity. Typically, mAb-enzyme conjugates are prepared using
15 bifunctional cross-linking reagents that react in a random fashion with exposed amino acid residues on the individual proteins. Immunoconjugates produced in this manner are microscopically heterogeneous due the inherent lack of specificity of the cross-linking reagents. In addition,
20 chemically prepared conjugates are typically isolated in low yields. Although recent reports describe alternative coupling chemistries that can afford higher yields of more homogeneous immunoconjugates (Mikolajczyk et al., Bioconjugate Chem.,
5:636-646 (1994); Werlen et al., Bioconjugate Chem., 5:411-417
25 (1994); Werlen et al., Tumor Targeting, 1:251-258 (1995)), these methods still involve chemical modification steps that can contribute to product heterogeneity.

Genetically constructed fusion proteins offer an alternative method of producing homogeneous mAb-enzyme
30 conjugates. There have now been reports describing the production, characterization, and activities of recombinant Fab, sFv, and disulfide stabilized Fv-enzyme fusion proteins (Bosslet et al., Br. J. Cancer, 65:234-238 (1992); Goshorn et al., Cancer Res., 53:2123-2127 (1993); Rodrigues et al.,
35 Cancer Res., 55:63-70 (1995)).

Thus, a need exists for homogeneous mAb-enzyme conjugates for use in combination with prodrugs. The instant invention addresses this need and more.

SUMMARY OF THE INVENTION

One aspect of the invention is a fusion polypeptide comprising antibody variable light and heavy region amino acid sequences specific for a melanoma associated antigen

5 operatively linked to a β -lactamase.

A further aspect of the invention is a method for the delivery of a cytotoxic agent to a tumor cell comprising the administration of a fusion polypeptide comprising antibody variable light and heavy region sequences operatively linked
10 to a β -lactamase, wherein the fusion polypeptide is reactive with a tumor cell antigen and the fusion polypeptide converts a prodrug to a cytotoxic drug, and the administration of an effective amount of the prodrug.

15 BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 depicts the structures of the cephalosporin mustard prodrug CCM and the parent drug phenylenediamine mustard PDM.

Fig. 2 depicts Scatchard binding analysis of L49
20 antibody binding to 3677 melanoma cells.

Fig. 3 depicts the construction of L49-sFv-bL. Three successive hybridization insertion reactions were used to install the 218 linker, variable heavy chain sequences, and variable light chain sequences into a pET phagemid containing
25 the r2-1 mutant of the *E. cloacae* β -lactamase. Single stranded phagemid DNA was produced by infection of XL-I Blue carrying the pET phagemids with M13KO7 helper phage. An oligonucleotide coding for the 218 linker sequence (- strand), with complementary regions to the 3' end of the pelB sequence
30 and the 3' end of the β -lactamase gene was prepared by chemical synthesis. Corresponding V_H and V_L sequences (- strand) were generated by asymmetric PCR.

Figs. 4A and 4B depicts the nucleotide (SEQ ID. NO: 19) and amino acid (SEQ ID. NO: 20) sequence for L49-sFv-bL
35 including the PelB leader sequence.

Figs. 5A and 5B depict SDS-PAGE (4-20%) and isoelectric focusing analyses (pH 3-10) of recombinant sp97 and wild type p97 antigen. Fig. 5A depicts non-reducing and

reducing SDS-PAGE analysis of recombinant sp97 and p97 antigen.

Figs. 6A-6D depict SDS-PAGE analyses of L49-sFv-bL expression and purification. Fig. 6A depicts the induction of L49-sFv-bL at varying IPTG concentrations (30°C, total cellular protein, 12% tris-glycine SDS-PAGE, Commassie staining, non-reducing conditions). The band corresponding to L49-sFv-bL is indicated with arrows. Fig. 6B depicts a Western analysis with rabbit polyclonal anti-bL. Lane 1: periplasm; Lane 2: L49-sFv-bL standard (12% tris-glycine SDS-PAGE, non-reducing conditions). Fig. 6C depicts the purification of L49-sFv-bL. Lane 1, periplasm; Lane 2, flow through from the sp97 affinity column; Lane 3, material that eluted from the sp97 column at pH 11; Lane 4, material that bound and eluted off the phenylboronic acid column (12% tris-glycine SDS-PAGE, Commassie staining, non-reducing conditions). Fig. 6D is a representative comparison of L49-sFv-bL to chemically prepared L49-Fab'-bL. Lane 1, L49-Fab'-bL; Lane 2, L49-sFv-bL (10% tris-glycine SDS-PAGE, Commassie staining, non-reducing conditions).

Fig. 7 depicts a competition binding assay. 3677 cells were incubated with various combinations of the test samples (L49, L49-sFv-bL, L49-Fab'-bL, and FITC-modified whole L49, keeping the total mAb concentration (test sample + L49-FITC) constant at 400 nM. Fluorescence intensity was determined by fluorescence activated cell sorter analysis.

Fig. 8 depicts cytotoxic effects of mAb-bL + CCM combinations on 3677 melanoma cells as determined by the incorporation of [³H]thymidine into DNA. 3677 cells were incubated with the mAb-bL conjugates, washed, and treated with CCM for 1 h. The effects were compared to cells treated with CCM or PDM for 1 h without prior conjugate exposure and to cells that were treated with saturating, amounts of unconjugated L49 prior to conjugate 1 treatment.

Figs. 9A and 9B depict pharmacokinetics of L49-sFv-bL in nude mice (3 animals/group). L49-sFv-bL was injected intravenously, tissues were removed and extracted at the indicated times, and the β -lactamase activity was determined

using nitrocefin as a substrate. Fig. 9A depicts clearance of L49-sFv-bL from the blood. Injected dose was 4 mg/kg. Fig. 9B depicts L49-sFv-bL levels in subcutaneous 3677 melanoma tumors and in normal tissues. Injected dose was 1 mg/kg.

5 Figs. 10A-10D. Therapeutic effects of L49-sFv-bL/CCM combinations in nude mice (six mice/group) with subcutaneous 3677 melanoma xenografts. Conjugates were injected, followed at various times by CCM (arrows on the X-axis). The average tumor volumes were reported until most or all of the animals were cured (tumors that became non-palpable for ≥ 10 tumor volume doubling times), or until an animal was removed from the experiment due to tumor outgrowth. Fig. 10A. L49-sFv-bL (1 mg/kg/injection) 12 h before CCM. Fig. 10B. L49-sFv-bL (1 mg/kg/injection) 24 h before CCM. Fig. 10C. L49-sFv-bL (4 mg/kg/injection) 24 h before CCM. Fig. 10D. L49-sFv-bL (4 mg/kg/injection) 48 h before CCM.

DETAILED DESCRIPTION OF THE INVENTION

The present invention encompasses pharmaceutical compositions, combinations and methods for treating cancers and other tumors. More particularly, the invention includes combinations comprising the antibody-enzyme fusion polypeptides of the invention and a corresponding prodrug or prodrugs for use in a method for treating tumors wherein a mammalian host is treated in a pharmaceutically acceptable manner with a pharmaceutically effective amount of an antibody-enzyme fusion polypeptide and a pharmaceutically effective amount of a prodrug or prodrugs. The combination and methods of this invention are useful in treating any mammal, including humans, dogs, cats, and horses.

In some embodiments the enzyme component of the antibody-enzyme fusion is a β -lactamase. The β -lactamase (bL) can be a class A, B, C, or D enzyme (Ledent et al., Biochem. J., 292:555-562 (1993); Felici et al., Biochem. J., 291:151-155 (1993)). Preferably, the bL is a class C enzyme. More preferably, the bL is an *Enterobacter cloacae* enzyme (Dubus et al., Biochem. J., 301:485-494 (1994)). The enzyme can be wild-type or mutant. For example, the *E. cloacae* P99 bL fusion

protein can be substituted at amino acid residues 537-541 (Gly Ser Asp Ser Lys (SEQ ID. NO: 1)). Possible substitutions at this region include Thr Ser Phe Gly Asn (SEQ ID. NO: 2), Ala Ser Ala Arg Arg (SEQ ID. NO: 3), Asn Asn Ala Gly Tyr (SEQ ID. NO: 4), Glu Val Glu Ile Lys (SEQ ID. NO: 5), Leu Thr Ser Asn Arg (SEQ ID. NO: 6), Gly Ser Lys Ser His (SEQ ID. NO: 7), Val Thy Arg Asn Gln (SEQ ID. NO: 8), Ile Val Asn Asn Lys (SEQ ID. NO: 9), Thr Ala Ile Pro Asp (SEQ ID. NO: 10) and Ile Thr Lys Pro Asp (SEQ ID. NO: 11) (Siemers et al., Biochemistry 35:2104-2111 (1996). A preferred mutant is Thr Ser Phe Gly Asn (SEQ ID. NO: 2).

The antibody component of the fusion can be obtained from any species. In an embodiment, the antibody is a murine monoclonal antibody directed against an antigen characteristic of a tumor cell to be targeted. Techniques for generating such monoclonal antibodies are well known in the art.

Antibodies can also be obtained from phage display and bacterial surface display libraries. This includes those produced from human tissue sources (to make human monoclonal antibodies) and combinatorial libraries of monoclonal antibodies produced by mutagenesis of CDR loops.

Typically, nucleic acid sequences encoding the variable regions of the light chains of the antibody are cloned by techniques well known in the art, such as by random hexamer primed reverse transcription reactions and PCR with signal sequence and constant region PCR primers (Jones et al., Biotechnology, 9:88-92 (1991)).

The corresponding DNA sequences are inserted into a vector of choice. Typically a secretory leader is operatively linked to the V_H and V_L region sequences. Any secretory leader can be used that will direct the secretion of the fusion protein into the periplasm of *E. coli* or other host, or into the medium. Exemplary secretory leaders include PelB, OmpA and StIII.

Typically the V_H and V_L region coding sequences are separated by a spacer that is generally greater than or equal to 15 amino acids. Examples include the sequence Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser (SEQ ID. NO:

12) and the sequence Gly Ser Thr Ser Gly Ser Gly Lys Pro Gly Ser Gly Glu Gly Ser Thr Lys Gly (SEQ ID. NO: 13) (218 linker). The V_L region gene is linked in continuous reading frame with the enzyme.

5 The entire fusion construct can be replicated as part of a vector that can be propagated in a host of interest, including but not limited to bacterial, yeast, insect, and mammalian hosts.

10 The prodrugs are preferably cephalosporin derivatives of phenylenediamine mustard, doxorubicin, mitomycin C, paclitaxel, vinca alkaloids, and melphalan. Other anti-cancer agents with active amines, hydroxyl or thiol groups can be modified with cephalosporins to form drugs that are activated by β -lactamases.

15 According to a preferred embodiment, the antibody-enzyme fusion polypeptide is administered prior to the introduction of the prodrug into the host. Sufficient time should be allowed between administration of the conjugate and the prodrug to allow the antibody component of the conjugate
20 to target and localize the enzyme to the tumor site. Such sufficient time may range from 4 hours to one week depending upon the conjugate used.

 The conjugates and prodrugs of the invention can be administered using conventional modes of administration
25 including, but not limited to, intravenous, intraperitoneal, oral, intralymphatic, or administration directly into the tumor. Intravenous administration is preferred.

 The compositions of the invention—comprising the immunoconjugates or prodrugs—may be in a variety of dosage
30 forms which include, but are not limited to, liquid solutions or suspensions, tablets, pills, powders, suppositories, polymeric microcapsules or microvesicles, liposomes, and injectable or infusible solutions. The preferred form depends upon the mode of administration and the therapeutic
35 application. For example, oral administration of the antibody-enzyme conjugate may be disfavored because the conjugate proteins tend to be degraded in the stomach if taken orally, e.g., in tablet form.

The conjugate or prodrug compositions also preferably include conventional pharmaceutically acceptable carriers and adjuvants known in the art such as human serum albumin, ion exchangers, alumina, lecithin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, and salts or electrolytes such as protamine sulfate.

The most effective mode of administration and dosage regimen for the compositions of this invention depends upon the severity and course of the disease, the patient's health and response to treatment and the judgement of the treating physician. Accordingly, the dosages of the immunoconjugates and prodrugs should be titrated to the individual patient.

Nevertheless, an effective dose of the antibody-enzyme conjugate of this invention may be in the range of from about 1.0 to about 100 mg/m². An effective dose of the prodrug of the invention will depend upon the particular prodrug used and the parent drug from which it is derived. Since the prodrug is less cytotoxic than the parent drug, dosages in excess of those recognized in the art for the parent drug may be used. For example, an effective dose of the cephalosporin mustard prodrugs may be in the range of about 500 mg/m².

In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting the scope of this invention in any manner.

EXPERIMENTAL EXAMPLES

Materials. The *Enterobacter cloacae* P99 gene was obtained from the plasmid pNU363 (Galleni et al., Biochem. J., 250:753-760 (1988)). The r2-1 bL was isolated from a library of *E. cloacae* enzymes in which the nucleotides corresponding to amino acids 286-290 were subjected to codon-based mutagenesis (Siemers et al., Biochemistry, 35:2104-2111 (1996)), and contains the residues Thr Ser Phe Gly Asn (SEQ ID. NO: 2) at these positions. L49-Fab'-bL was prepared as previously described by combining thiolcontaining Fab' fragments of the antibodies with maleimide-substituted bL,

forming a thioether link between the two proteins (Svensson et al., Bioconjugate Chem., 5:262-267 (1994)).

Isolation and Characterization of the L49 Antibody.

The L49 producing hybridoma was developed using standard techniques as previously described for the isolation of other hybridomas (Yeh et al., Proc. Natl. Acad. Sci. (USA), 76:2927-2931 (1979)). Balb/C mice were immunized repeatedly with the H2981 (lung carcinoma), CH3 (lung carcinoma), and W56 (melanoma) cell lines, all of which were derived from human tumors. Spleen cells from the immunized mice were hybridized with the neomycin gene transfected myeloma cell line P3X63-Ag8.563 (Yeh et al., Proc. Natl. Acad. Sci. (USA), 76:2927-2931 (1979)). Standard selection and cloning yielded a hybridoma producing the L49 IgG1 antibody.

Scatchard analysis of L49 binding was performed by radiolabeling the mAb with [¹²⁵I]Iodogen to a specific activity of 0.3 mCi/mg protein. 3677 melanoma cells (Kerr et al., Cancer Res., 55:3558-3563 (1995)) in 96-well plates (13,000 cells/well) were incubated with 0.03 to 10 nM ¹²⁵I-L49 for 30 min. on ice, and then the cells were separated from unbound radioactivity by centrifugation through silicon oil. The tubes were frozen, the cell pellet was cut from the supernatant, and both fractions were counted in a gamma counter (Packard). Binding affinity and sites per cell were determined by Scatchard analysis (Trucco and dePetris, Immunological Methods, I. Lefkovits and B. Pernis, eds., Academic Press, New York, pp. 1-26 (1981)).

Soluble p97 (sp97). A secreted form of p97 (sp97) was made utilizing PCR based mutagenesis to introduce a stop codon at the cysteine residue three amino acids upstream of the glycoposphatidylinositol anchor domain (Alemany et al., J. Cell Sci., 104:1155-1162 (1993); Food et al., J. Biol. Chem., 269:3034-3040 (1994)). Briefly, the 3' oligonucleotide used in the PCR reaction contained the mutation changing the S710 codon to a stop codon. In all, the coding sequence for 29 amino acids was deleted from the carboxyl terminus of wild type p97. Cloning and expression of sp97 was accomplished using a glutamine synthetase gene as an amplifiable marker in

CHO cells (Cockett et al., Biotechnology, 8:662-667 (1990)). The sp97 gene was cloned into pEE14 (Stephens and Cockett, Nucl. Acids Res., 17:7110 (1989)) and transfected into CHO-K1 cells by calcium phosphate coprecipitation. Transformants were initially selected for resistance to 25 μ M methionine sulfoximine, and sp97 secreting colonies were selected for amplification at drug concentrations of 100, 250, and 500 μ M. The selection and amplification media used was Glasgow Minimum Essential Medium without L-glutamine, tryptose phosphate broth or sodium bicarbonate supplemented with 10% dialyzed fetal bovine serum. A cloned CHO cell line secreting sp97 was cultured in 10-shelf cell factories containing 1.5 liters of media.

Soluble p97 was isolated on a 96.5 immunoaffinity chromatography column as described for the purification of wild type p97 from melanoma cells (Baker et al., FEBS Lett., 298:215-218 (1992)). Small amounts of residual contaminants were removed by gel filtration on a Sephacryl S300 HR column (Pharmacia LKB) using PBS as eluant. Solutions containing sp97 were concentrated by ultrafiltration to 1-5 mg/ml, sterilized by passage through a 0.1 μ filter, and stored at 2-8°C for up to 6 months without noticeable loss of biochemical or biological activity.

Cloning of L49 Variable Regions and sFv Construction.

Construction of L49-sFv-bL by hybridization insertion was performed with materials and protocols from the BioRad M13 mutagenesis kit, except for isolation of single stranded phagemid template (Qiagen M13 kit, M13KO7 helper phage). The variable regions of the L49 antibody were cloned from the corresponding hybridoma mRNA by reverse transcription-PCR (Perkin Elmer GeneAmp reagents and Model 9600 thermal cycler), using random hexamer primed reverse transcription reactions and signal sequence and constant region PCR primers (Jones et al., Biotechnology, 9:88-92 (1991)). Construction of L49-sFv-bL began with a single-stranded template of the pET-26b phagemid containing the r2-1 mutant of the *Enterobacter cloacae* P99 bL gene (Siemers et al., Biochemistry, 35:2104-2111 (1996)) fused to the pelB leader sequence. Hybridization

mutagenesis was used to insert the 218 linker sequence (Whitlow et al., Protein Eng., 6:989-995 (1993)) (chemically synthesized oligonucleotide, 5'- TTCTGACACTGGCGTGCCCTTGGTA GAGCCTTCGCCAGAGCCCGGTTTGCCAGAGCCGGACGTCGAGCCGGCCATCGCCGGCTG-3' (SEQ ID. NO: 14)) and full V_H and V_L region sequences (oligonucleotides produced by asymmetric PCR, V_H forward primer: 5'-CCAGCCGGCGATGGCCGAGGTGCAGCTTCAGGAGT-3' (SEQ ID. NO: 15); V_H reverse primer: 5'-AGAGCCGGACGTCGAGCCTGAGGAGACGGT GACAGAGG-3' (SEQ ID. NO: 16); V_L forward primer: 5'- AGGCTCTACCAAGGGCGATTTTGTGATGACCCAAAC-3' (SEQ ID. NO: 17); V_L reverse primer: 5'-TTCTGACACTGGCGTCCGTTTGATTTCAGCTTGG-3' (SEQ ID. NO: 18) in between the pelB leader sequence and bL in a 5'-pelB-V_H-218-V_L-bL-3' orientation. The nucleotide (SEQ ID NO: 19) and amino acid (SEQ ID NO. 20) sequences of L49-sFv-bL are provided in Figs. 4A and 4B.

Expression, Purification and Characterization of L49-sFv-bL. L49-sFv-bL was expressed as a soluble protein in *E. coli* strain BL21 (λ DE3) at 23°C in 4L, baffled shake flasks. T-broth (1L) containing 30 μ g/ml kanamycin was inoculated with several colonies of freshly transformed BL21 (λ DE3) cells. The flasks were shaken (200 rpm) at 37°C until the absorbance at 660 nm reached 0.8. The culture was cooled to 23°C and IPTG (50 μ M) was added. The culture was incubated with shaking for an additional 16 h at 23°C, at which time the absorbance at 660 nm was between 8-15. The cells were pelleted by centrifugation and resuspended in 30 mM Tris, 2 mM EDTA, 0.3% (v/v) Nonidet P-40, pH 8.5, 4°C. The mixture was stirred gently for 1 h, repelleted, and the supernatant was decanted and filtered (0.2 μ m).

Purification of L49-sFv-bL was accomplished by a two-step affinity purification. The periplasmic fraction was first applied to a Sepharose column of immobilized sp97 antigen. The column was washed with PBS until the absorbance at 280 nm reached the baseline level, and bound protein was eluted with pH 11.2 buffer (50 mM sodium phosphate, 100 mM NaCl). Fractions containing the bound protein were neutralized with 1/10 v/v of 3M phosphate, pH 7.2. This material was then subjected to Sepharose 4B m-

aminophenylboronic acid affinity chromatography (Cartwright et al., Biochem. J., 221:505-512 (1984)) using washing and elution conditions described above. The resulting preparation was dialyzed against PBS, filtered (0.2 μ m), and stored at 4°C (0.1-1.1 mg/ml).

Competition binding experiments were performed as described by Svensson et al. (Bioconjugate Chem., 3:176-181 (1992)). Immunoassays were performed by coating polystyrene 96 well plates with sp97 (0.1 mL, 2 μ g/ml in PBS, overnight, 4°C.) After blocking with specimen diluent (Genetic Systems Corp.) for 1 h at 22°C, the blocking solution was replaced with fresh specimen diluent (0.1 ml) containing serial dilutions of the samples. After 1 h at 22°C the plates were washed, followed by development with 0.1 mL of a nitrocefin (O'Callaghan et al., Antimicrobial Agents and Chemotherapy, 1:283-288 (1972)) solution (0.1 mM in PBS/1% dimethylformamide, 15 min, 22°C. Absorbance measurements were read in an plate reader using a 490 nm filter with 630 nm as the reference wavelength.

In Vitro Cytotoxicity. 3677 melanoma cells were plated into 96-well microliter plates (10^4 cells/well in 100 μ L of Iscove's Modified Dulbecco's Medium (IMDM) with 10% fetal bovine serum, penicillin (60 μ g/ml), and streptomycin (100 μ g/ml) and allowed to adhere overnight. For blocking experiments, the cells were incubated with unconjugated L49 at 1 μ M for 30 min. prior to treatment with the L49 conjugates. The cells were treated with L49-sFv-bL or L49 Fab'-bL at 10 nM. After 30 min. at 4°C, the plates were washed three times with antibiotic-free RPMI 1640 media (Gibco) with 10% fetal bovine serum, and then varying concentrations of CCM (Fig. 1) were added. CCM and PDM were also added to cells treated with media alone. After 1 h at 37°C, cells were washed three times with IMDM and incubated approximately 18 h at 37°C. The cells were then pulsed for 12 h with [3 H] thymidine (1 μ Ci/well) at 37°C, detached by freezing at -20°C and thawing, and harvested onto glass fiber filter mats using a 96-well harvester (Wallac, Gaithersburg, MD). Radioactivity was counted using a LKB Wallac β -plate counter.

In Vitro Cytotoxicity. 3677 melanoma cells were plated in 96-well microtiter plates (10^4 cells/well in 100 μ L of IMDM with 10% fetal bovine serum, 60 μ g/ml penicillin and 0.1 mg/ml streptomycin) and allowed to adhere overnight. For blocking experiments, the cells were incubated with unconjugated L49 at 1 μ M for 30 min prior to treatment with the L49 conjugates. The cells were treated with L49-sFv-bL or L49 Fab'-bL at 10 nM. After 30 min at 4°C, the plates were washed three times with antibiotic-free RPMI 1640 media (Gibco) with 10% fetal bovine serum, and then varying concentrations of CCM were added. CCM and PDM were also added to cells treated with media alone. After 1 h at 37°C, cells were washed three times with IMDM and incubated approximately 18 h at 37°C. The cells were then pulsed for 12 h with [3 H] thymidine (1 μ Ci/well) at 37°C, detached by freezing at -20°C and thawing, and harvested onto glass fiber filter mats using a 96-well harvester. Radioactivity was counted using a LKB Wallac β -plate counter.

Conjugate Localization. Subcutaneous 3677 melanoma tumors were established in female athymic nu/nu mice (8-12 weeks old, Harlan Sprague-Dawley, Indianapolis, IN) by transplanting tumors that had been previously passaged as previously described (Kerr et al., Cancer Res., 55:3558-3563 (1995)). Tumor bearing mice were injected i.v. with L49-sFv-bL (1 or 4 mg mAb component/kg) or with L49-Fab'-bL (1.8 mg mAb component/kg). At various time intervals, the mice were anesthetized, bled through the orbital plexus, and sacrificed. Tissues were removed and homogenized in PBS containing 15 μ g/ml aprotinin (2 mL/g tissue). To the homogenate was added 50 mM sodium phosphate containing 100 mM NaCl at pH 11.2 (10 mL/g tissue), and the suspension was mixed. After 20 min at room temperature, 3M sodium phosphate at pH 7.0 was added (2 mL/g tissue), and the mixture was mixed and centrifuged.

Quantification of conjugate concentrations was accomplished using a direct enzyme immunoassay. Polystyrene 96-well microtiter plates were coated with an affinity-purified rabbit polyclonal antisera to wild type *E. cloacae* bL (1 μ g/ml), and were then blocked with specimen diluent

(Genetic Systems Corp.). Serially diluted tissue extracts or purified samples (L49-sFv-bL as a standard for the fusion protein samples, L49-Fab'-bL as a standard for the L49-Fab'-bL samples) were added to the wells and allowed to bind for 3 h at room temperature. The plates were washed and developed by the addition of 0.1 ml of nitrocefin (O'Callaghan et al., Antimicrobial Agents and Chemotherapy, 1:283-288 (1972)) at 0.1 mM in PBS containing 1% dimethylformamide. Absorbance measurements were read in an ELISA plate reader using a 490 nm filter with 630 nm as the reference wavelength.

In Vivo Therapy Experiments. 3677 tumor-bearing mice (subcutaneous implants, six animals/group, average tumor volume 130 mm³) were injected with L49-sFv-bL (i.v., 7-8 days post tumor implant), followed 12-48 h later by CCM using doses of fusion protein and prodrug as indicated in the Results section. Treatment with L49-sFv-bL + CCM was repeated 1 week later. Animals were monitored 1-2 times/week for body weight, general health, and tumor growth. Tumor volume was estimated using the formula: longest length X perpendicular dimension²/2. Cures were defined as an established tumor that, after treatment, was not palpable for ≥ 10 tumor volume doubling delays (≥ 40 days in the 3677 tumor model). Maximum tolerated doses led to less than 20% weight loss, no treatment-related deaths, and were within 50% of the dose where such events took place.

RESULTS

Characterization of the L49 Antibody. The L49 antibody (IgG1) binds to the p97 antigen, which has been shown to be present on most human melanomas and many carcinomas (Brown et al., J. Immun., 127:539-545 (1981), Brown et al., Proc. Natl. Acad. Sci. (USA), 78:539-543 (1981); Woodbury et al., Proc. Natl. Acad. Sci. (USA), 77:2183-2187 (1980)). Scatchard analysis of the binding of radiolabeled L49 to the 3677 human melanoma cell line indicated that the mAb bound with a dissociation constant of 1.0 nM (Fig. 2). At saturation, there were approximately 2.1×10^4 molecules of L49 bound/cell. These values are very similar to those

obtained for the 96.5 mAb (Kerr et al., Cancer Res. 55:3558-3563 (1995)), which also binds to p97, but to a different epitope than L49 (data not shown).

5 **Cloning and Expression of L49-sFv-bL.** The variable region genes for the L49 antibody heavy and light chains were cloned from the L49 hybridoma line by RT-PCR of hybridoma mRNA and amplification of the corresponding cDNA. A consensus sequence was determined by examining several clones from independent reverse transcription reactions to reduce the
10 possibility of reverse transcription or PCR derived errors. The PCR primers used were complementary to the signal sequence and constant region of the mAb. Thus, the entire variable regions were obtained.

15 L49-sFv-bL was constructed in a stepwise fashion by hybridization insertion of the sFv linker, V_H , and V_L region sequences onto a single stranded pET phagemid template containing the pelB leader sequence and bL gene (Fig. 3). The particular bL gene used encoded a mutated form of bL (r2-1) that contained the sequence Thr Ser Phe Gly Asn (SEQ ID. NO: 2) at positions 286-290. This mutated bL has been shown to
20 have slightly greater activity than the wild type enzyme (Siemers et al., Biochemistry, 35:2104-2111 (1996)). The 218 linker sequence corresponds to amino acids Gly Ser Thr Ser Gly Ser Gly Lys Pro Gly Ser Gly Glu Gly Ser Thr Lys Gly (SEQ ID. NO: 13), and was used as the sFv linker based on its ability
25 to reduce sFv protein aggregation (Whitlow et al., Protein Eng., 6:989-995 (1993)). An oligonucleotide coding for the 218 linker (- strand, produced by chemical synthesis) was first annealed to the phagemid template, resulting in a pelB-218-bL construct. V_H and V_L region segments (produced by
30 asymmetric PCR) were then inserted into the intermediate construct to generate the final L49-sFv-bL gene in an pelB- V_H -218- V_L -bL orientation. The pelB leader sequence results in transport of the protein into the periplasmic space of *E.*
35 *coli*. No additional linker was placed between V_L and the bL enzyme. The nucleotide (SEQ ID NO: 19) and amino acid (SEQ ID NO: 20) sequences are provided in Figs. 4A and 4B.

To facilitate the isolation and characterization of L49-containing fusion proteins, a soluble form of the p97 antigen was developed. This was made by truncating the p97 antigen at a site upstream to the membrane anchoring domain.

5 The soluble antigen (sp97) was expressed in CHO-K1 cells and purified by affinity chromatography. SDS-PAGE analysis of recombinant sp97 indicated that it was slightly lower in molecular weight than p97 (Fig. 5A). Isoelectric focussing revealed little difference between p97 and sp97 (Fig. 5B), a

10 result that was anticipated, since only a single charged residue is lost in the sp97 construct.

L49-sFv-bL was expressed in soluble form in an *E. coli* strain that was transformed with the plasmid shown in Fig. 3. Quantitation of L49-sFv-bL containing samples was performed

15 using an immunoassay in which the L49 portion was captured onto microtiter plates that were coated with sp97, and the bL enzyme activity was determined using nitrocefin as a calorimetric indicator for bL activity (O'Callaghan et al., Antimicrobial Agents and Chemotherapy, 1:283-288 (1972)).

20 Thus, only bifunctional fusion protein was measured. Under the transcriptional control of the T7 promoter and lac operon, fusion protein expression could be detected by SDS-PAGE analyses of cell pellets when at IPTG concentrations as low as 1.6 μ M (Fig. 6A). Significant levels of toxicity were

25 observed when the IPTG concentration exceeded 90 μ M, resulting in inhibition of cell growth and in the eventual outgrowth of cell populations that did not express fusion protein. Typically, 50 μ M IPTG induction was used for large scale experiments, since this led to higher levels of fusion protein

30 expression without significant levels of cytotoxicity. It was also found that expression of L49-sFv-bL was enhanced at lower temperatures, such that protein yields were higher at 23 or 30°C compared to 37°C. Similar results have been noted for the expression of antibody fragments and other recombinant

35 proteins in *E. coli* (Plückthun, Immun. Rev., 130:151-188 (1992)).

In shake flask cultures, 80% of active material was present in the periplasm of the bacterial cells, with the

remainder present in the culture supernatant. Conventional techniques for releasing the periplasmic contents, such as sucrose/lysozyme spheroplasting or osmotic shock, resulted in only a limited release of the available protein. Similar results were also obtained using 20% alone. Freeze-thawing or sonication of cells to release total cytoplasmic material did not result in an increased yield of functional fusion protein. It was found that a high yield of fusion protein could be obtained by treating cell pellets with the detergent Nonidet-P-40. Expression levels of L49-sFv-bL using this detergent ranged from 2.5-8 mg/L culture. Western analysis (Fig. 6B) with a rabbit polyclonal antisera raised to bL showed that most of the bL containing protein in the preparation was approximately 63 kDa in molecular weight (theoretical molecular weight 66.5 kDa). Small amounts of truncated fragments and aggregated material were also detected.

Purification of L49-sFv-bL. The purification of L49-sFv-bL to homogeneity was achieved by a two-step affinity chromatography procedure. Periplasmic preparations from shake flask cultures were first applied to an immobilized sp97 affinity column that was capable of binding to the L49 portion of the conjugate. After extensive washing, bound material was eluted at pH 11.2 (Fig. 6C). Acidic pH conditions (pH 2.2) successfully eluted the fusion protein but caused precipitation of material when working with multi-milligram quantities of fusion protein. The sp97 chromatography purified material was approximately 70% pure by size-exclusion HPLC and SDS-PAGE, with the contaminants consisting of two bands of approximately 33 kDa molecular weight. The second step of the purification involved binding the material to immobilized phenylboronic acid. This resin has previously been found to bind to β -lactamases, presumably to the active site of the enzymes (Cartwright and Waley, Biochem. J. 221;505-512 (1984)). This led to the recovery of protein that was pure by SDS-PAGE analysis (Figs. 6C and 6D).

L49-sFv-bL Characterization and Activity. In view of the detergent based release of L49-sFv-bL from the bacteria, it was important to demonstrate that isolated fusion protein

had been correctly processed and transported into the periplasm, such that the pelB leader sequence was cleaved from the amino terminus of the V_H region. This was determined by
5 analysis. The sequence obtained (Glu Val Leu Gln Leu Glu Ser (SEQ ID. NO: 21)) was identical to the expected V_H amino terminal sequence, indicating that the leader sequence was proteolytically clipped, as designed.

10 The binding characteristics of the sFv portion of the fusion protein were determined using a fluorescent activated cell sorting competition assay in which fusion protein and FITC-modified whole L49 competed for binding to cell-surface antigens on SK-MEL 28 melanoma cells. L49-sFv-bL and the L49
15 Fab'-bL chemical conjugates bound equally well to the cell line, indicating that the binding activity of the antibody portion of the conjugate was preserved (Fig. 7). More detailed information about binding was obtained using surface plasmon resonance, which allowed the measurement of the on and off rates of L49-sFv-bL binding to the p97 antigen immobilized
20 on a gold surface (Table 1). This assay established that the binding affinity of the fusion protein to the p97 antigen (K_d = 1.0 nM) was comparable to L49 Fab' (K_d = 0.73 nM), and chemically produced L49-Fab'-bL conjugate (K_d = 1.3 nM).

Table 1. Binding and enzyme kinetic parameters of L49 and bL containing proteins

Sample	$k_{on}(M^{-1} \cdot s^{-1})$	$k_{off}(s^{-1})$	$k_D(nM)$	$k_{cat}(s^{-1})$	$k_m(\mu M)$
L49 Fab'	2.3×10^5	1.7×10^{-4}	0.73	n.a. ^b	n.a.
r2-1 bL ^c	n.a.	n.a.	n.a.	261	19
L49-Fab'-bL ^d	1.8×10^5	2.4×10^{-4}	1.3	n.d. ^e	n.d.
L49-sFv'-bL ^c	4.1×10^5	4.2×10^{-4}	1.0	232	19

19

^a Values shown are the average of a minimum of two independent experiments, except for L49-Fab'-bL (binding experiment performed once). The range of values obtained in Michaelis-Menten kinetic analyses were within 5% of the means.

^b Not Applicable.

^c The r2-1 bL contains mutations at positions 286-290 compared to the wild type enzyme (Siemers et al., Biochemistry 35:2104-2111 (1996)).

^d Chemically prepared conjugate containing the wild type enzyme.

^e Not determined.

Enzymatic activity assays of the bL portion of L49-sFv-bL were undertaken using nitrocefin as the enzyme substrate (Table 1). Michaelis-Menten kinetic analyses confirmed that the fusion protein retained the full enzymatic activity of the mutant bL, enzyme from which it was derived (Siemers et al., Biochemistry, 35:2104-2111 (1996)). Thus, both the binding of the L49 antibody and enzymatic activity of the *E. cloacae* r2-1 bL were preserved in the fusion protein.

The cytotoxic effects of L49-sFv-bL in combination with CCM were determined on 3677 human melanoma cells, which express the p97 antigen. The experiments were performed by exposing the cells to the conjugates, and washing off unbound material before adding various concentrations of CCM. [³H]-Thymidine incorporation was used to measure cytotoxic activity. The prodrug CCM (IC₅₀, 16 μM) was approximately 50-fold less toxic to 3677 cells than PDM (IC₅₀, 0.3 μM). As expected, L49-sFv-bL and L49-Fab'-bL were equally effective at prodrug activation, and the combinations were equivalent in activity to PDM (Fig. 8). This indicates that prodrug conversion was efficient under the conditions tested. In addition, it was found that activation was immunologically specific, since L49-sFv-bL did not activate CCM on cells that were previously saturated with L49 mAb before being exposed to the fusion protein.

In Vivo Localization. Biodistribution studies of L49-sFv-bL and L49-Fab'-bL were carried out in nude mice bearing subcutaneous (s.c.) 3677 melanoma tumor xenografts. The conjugates were injected i.v., and at various time points tissues were removed and extracted under alkaline conditions to disrupt antigen-antibody interactions. The samples were then trapped a polyclonal antiserum to bL, and bL activity was measured using nitrocefin as a colorimetric indicator (O'Callaghan et al., Antimicrobial Agents and Chemotherapy, 1:283-288 (1972)). Control experiments in which L49-sFv-bL was directly injected into excised tumors and tissues indicated that this extraction procedure recovered 90% of the injected bL activity.

L49-sFv-bL cleared very rapidly from the blood (Fig. 9A). The initial and terminal clearance half lives ($t_{1/2\alpha}$ and $t_{1/2\beta}$) were 0.3 and 2.5 h, respectively, leading to a 10^4 reduction of L49-sFv-bL blood levels within 24 h of conjugate administration. In spite of this rapid clearance, relatively high intratumoral levels of L49-sFv-bL were measured compared to normal tissues, and the ratio remained high for 24 h (Fig. 9B). At 4 h post L49-sFv-bL administration, the tumor to blood ratio was 13:1. The ratio increased substantially with time, and was 105:1 within 24 h of conjugate administration (Table 2).

Table 2. Tissue Distribution of Immunoconjugates

Treatment ^a	Time	% Injected dose/g (standard deviation)				
		Tumor	Liver	Spleen	Kidney	Blood
L49-sFv-bL, 1 mg/kg tumor/tissue ratios	4 hrs	1.1 (0.2) 1	0.021 (0.002) 52	0.014 (0.008) 79	0.027 (0.015) 41	0.084 (0.04) 13
L49-sFv-bL, 1 mg/kg tumor/tissue ratios	12 hrs	0.53 (0.17) 1	<0.003 >177	<0.003 >177	<0.003 >177	0.008 (0.001) 66
L49-sFv-bL, 1 mg/kg tumor/tissue ratios	24 hrs	0.21 (0.01) 1	<0.003 >70	<0.003 >70	<0.003 >70	0.002 (0.001) 105
L49-sFv-bL, 4 mg/kg tumor/tissue ratios	12 hrs	0.73 (0.02) 1	<0.003 >240	<0.003 >240	<0.003 >240	0.009 (0.001) 81
L49-sFv-bL, 4 mg/kg tumor/tissue ratios	24 hrs	0.29 (0.05) 1	<0.001 >290	<0.001 >290	<0.001 >290	0.002 (0.0002) 141
L49-sFv-bL, 4 mg/kg tumor/tissue ratios	48 hrs	0.15 (0.07) 1	n.d. ^b	n.d.	n.d.	0.001 (0.0002) 150
L49-Fab'-bL, 1.8 mg/kg tumor/tissue ratios	72 hrs	0.28 (0.26) 1	0.015 (0.003) 19	0.010 (0.006) 28	0.016 (0.005) 18	0.05 (0.015) 5.6

^a Mice (3 animals/group) were injected with conjugates, and at the times indicated, tissues were excised and extracted to remove the conjugate. The percent injected dose was based on the measured bL activity compared to standard curves obtained from extracted tissues that were spiked with known amounts of L49-sFv-bL or L49-Fab'-bL.

^b Not determined.

Similar results were obtained using L49-sFv-bL doses of 4 mg/kg. At this dose, very high tumor to blood ratios (141-151:1) were measured 24-48 h after the conjugate was administered. Interestingly, chemically produced L49-Fab'-bL cleared quite slowly from the blood and had only a 5.6:1 tumor to blood ratio 72 h after administration. Thus, L49-sFv-bL localizes in tumors, clears rapidly from the systemic circulation, and has significantly improved pharmacokinetic properties compared to the chemically produced L49-Fab'-bL conjugate.

Therapeutic Activity. In vivo therapy experiments were performed using the L49-sFv-bL/CCM combination in nude mice with established s.c. 3677 tumors. This particular tumor model has previously been shown to be resistant to treatment with doxorubicin, PDM, and CCM (Kerr et al., Cancer Res., 55:3558-3563 (1995)). In the experiments reported here, conjugate treatment was initiated 7-8 days after tumor implant, at which time the tumors were approximately 130 mm³ in volume. CCM was then administered 12, 24, or 48 h later, and the treatment protocol was repeated after 1 week. Maximum tolerated doses are defined as those that led to less than 20% weight loss, no treatment-related deaths, and were within 50% of the dose where such events took place. A tumor was considered as having been cured once it was not palpable for at least 10 tumor volume doubling times, based on the tumor growth of untreated animals (tumor volume doubling time was 4 days). If an animal was removed from the experiment because of tumor growth, the data from the entire group was no longer plotted, but the remaining animals were followed for tumor size and general health.

The maximum tolerated doses of CCM (300 mg/kg/injection) and PDM (3 mg/kg/injection) when administered weekly for three rounds induced 2 and 6 day delays in tumor outgrowth, respectively (data not shown). In contrast, pronounced antitumor activity was obtained in mice that received L49-sFv-bL prior to treatment with CCM (Fig. 10A). Therapeutic efficacy was schedule and dose dependent. Tumor cures were obtained in all of the animals that received CCM

(125 and 175 mg/kg/injection) 12 h after treatment with L49-sFv-bL (Fig. 10A). In this dosing schedule, significant antitumor activity including four cures in the group of six mice was obtained when the CCM dose was reduced to 75 mg/kg/injection. The remaining two animals in this group had tumors that underwent partial regressions, but eventually began to grow after the last prodrug treatment. There were no apparent toxicities in any of these treatment groups.

Significant antitumor activity could also be achieved when the prodrug was administered 24 h post conjugate administration, either by increasing the prodrug dose and keeping the conjugate dose constant at 1 mg/kg/injection (Fig. 10B), or by increasing the conjugate dose to 4 mg/kg/injection (Fig. 10C). In both cases, the majority of tumors were cured, again with no evidence of toxicity. Finally, therapeutic efficacy was also obtained with a 48 h interval between conjugate and prodrug administration (Fig. 10D). Tumor regressions were obtained in all of the mice in these groups, and three of six animals that received 275 mg/kg/injection CCM were cured. Thus, the antitumor activities of L49-sFv-bL in combination with CCM were pronounced, and therapeutic efficacy was achieved in a variety of dosing schedules.

In summary, the fusion protein L49-sFv-bL was constructed and expressed in *E. coli* as a soluble protein, and isolated from the periplasmic space using a two-stage affinity chromatography method. After purification, L49-sFv-bL was demonstrated to be homogeneous by SDS-PAGE, and was fully active with respect to both the L49 and bL components. As expected, the fusion protein was able to bind to melanoma cells that expressed the p97 antigen, and were able to activate a cephalosporin mustard in an immunologically specific manner.

To minimize systemic, non-targeted drug release *in vivo*, a high mAb-enzyme tumor to normal tissue ratio is needed before the anticancer prodrug is administered. To attain the required localization index in mice, the time between conjugate and prodrug administration has varied significantly from one conjugate to another. For example, the delay between

conjugate and prodrug administration was 3 days for 96.5-Fab'-bL (molecular weight 92 kDa, Kerr et al., Cancer Res., 55:3558-3563 (1995)), 1 week for the anti CEA-Fab- β -glucuronidase fusion protein (molecular weight 250 kDa, Bosslet et al., Cancer Res., 54:2151-2159 (1994)), and 2 weeks for the ICR12-carboxypeptidase G2 conjugate (molecular weight range of 233-316 kDa, Eccles et al., Cancer Res., 54:5171-5177 (1994)). In some cases, it has even been necessary to accelerate systemic conjugate clearance in a separate step involving the formation of immune complexes before prodrug could be administered (Wallace et al., Cancer Res., 54:2719-2723 (1994); Kerr et al., Bioconjugate Chem., 4:353-357 (1993); Rogers et al., Br. J. Cancer, 72:1357-1363 (1995)). Here L49-sFv-bL not only clears very rapidly from the systemic circulation (Fig. 9A), but also preferentially localized into subcutaneous tumor xenografts (Fig. 9B, Table 2). The very high tumor to non-tumor fusion protein ratios obtained within 4-12 hours of conjugate treatment would lead to the prediction that, in contrast to other enzyme/prodrug systems (Bosslet et al., Cancer Res., 54:2151-2159 (1994); Eccles et al., Cancer Res., 54:5171-5177 (1994); Kerr et al., Cancer Res., 55:3558-3563 (1995)), therapeutic efficacy would not require protracted time intervals between conjugate and prodrug administration. This has now been experimentally confirmed, since cures of established tumors were obtained when CCM was administered 12 h following the conjugate (Fig. 10A).

It is noteworthy that a correlation can be made between the outcome in the therapy experiments (Figs. 10A-D) and the pharmacokinetic data (Figs. 9A and B and Table 2). At a given conjugate dose, the intratumoral concentration decreased with a half life of approximately 8 h (Table 2). This may be due to a variety of factors such as dissociation of the conjugate from the antigen, membrane recycling, enzyme metabolism, and rapid tumor growth. The net result is that longer time intervals between conjugate and prodrug administration require that either the amount of prodrug or conjugate be increased in order to maintain therapeutic efficacy (Figs. 10A-D).

All references cited herein are specifically incorporated by reference in their entirety for all purposes.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, those with ordinary skill in the art will appreciate that other embodiments and variations of the invention are possible which employ the same inventive concepts described above. Therefore, the invention is not to be limited by the above disclosure, but is to be determined in scope by the claims which follow.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT:
 - (A) NAME: Bristol-Myers Squibb Company
 - (B) STREET: 345 Park Avenue
 - (C) CITY: New York
 - (D) STATE: New York
 - (E) COUNTRY: United States of America
 - (F) POSTAL CODE (ZIP): 10154
 - (G) TELEPHONE:
 - (H) TELEFAX:
 - (I) TELEX:
- (ii) TITLE OF INVENTION: RECOMBINANT ANTIBODY-ENZYME FUSION PROTEINS
- (iii) NUMBER OF SEQUENCES: 21
- (iv) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (v) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: WO
 - (B) FILING DATE: 30-APR-1998
- (vi) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 60/045,888
 - (B) FILING DATE: 07-MAY-1997
- (vii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Poor, Brian W.
 - (B) REGISTRATION NUMBER: 32,928
 - (C) REFERENCE/DOCKET NUMBER: 9197A-83-1PC
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (206) 467-9600
 - (B) TELEFAX: (415) 576-0300

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:
Gly Ser Asp Ser Lys
1 5

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Thr Ser Phe Gly Asn
1 5

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Ala Ser Ala Arg Arg
1 5

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Asn Asn Ala Gly Tyr
1 5

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Glu Val Glu Ile Lys
1 5

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Leu Thr Ser Asn Arg
1 5

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Gly Ser Lys Ser His
1 5

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Val Thr Arg Asn Gln
1 5

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Ile Val Asn Asn Lys
1 5

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Thr Ala Ile Pro Asp
1 5

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 5 amino acids
 - (B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Ile Thr Lys Pro Asp
1 5

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 16 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Gly Ser Thr Ser Gly Ser Gly Lys Gly Ser Gly Gly Ser Thr Lys Gly
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 84 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TTCTGACACT GCGTGCCCT TGGTAGAGCC TTCGCCAGAG CCCGGTTTGC CAGAGCCGGA 60
CGTCGAGCCG GCCATCGCCG GCTG 84

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

31

- (A) LENGTH: 35 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CCAGCCGGCG ATGGCCGAGG TGCAGCTTCA GGAGT

35

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

AGAGCCGGAC GTCGAGCCTG AGGAGACGGT GACAGAGG

38

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

AGGCTCTACC AAGGGCGATT TTGTGATGAC CCAAAC

36

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

TTCTGACACT GCGTCCGTT TGATTTCAG CTTGG

35

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2010 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

AATTAATACG ACTCACTATA GGGGAATTGT GAGCGGATAA CAATTCCCCT CTAGAAATAA	60
TTTGTGTTAA CTTTAAAGAAG GAGATATACA TATGAAATAC CTGCTGCCGA CCGCTGCTGC	120
TGGTCTGCTG CTCCTCGCTG CCCAGCCGGC GATGGCCGAG GTGCAGCTTC AGGAGTCAGG	180
ACCTAGCCTC GTGAAACCTT CTCAGACTCT GTCCCTCACC TGTTCGTCA CTGGCGACTC	240
CATCACCAGT GGTACTGGA ACTGGATCCG GAAGTTCCCA GGGAAATAAC TTGAATATAT	300
GGGTACATA AGCGACAGTG GTATCACTTA CTACAATCCA TCTCTCAAAA GTCGCATTTT	360
CATCACTCGA GACACATCCA AGAACCAATA CTACCTCCAG TTGAATTTTG TGAAGTCTGA	420
GGACACAGCC ACATATAACT GTGCAAGAAG GACTCTGGCT ACTTACTATG CTATGGACTA	480
CTGGGGTCAA GGAACCTCTG TCACCGTCTC CTCAGGCTCG ACGTCCGGCT CTGGCAAACC	540
GGGCTCTGGC GAAGGCTCTA CCAAGGGCGA TTTTGTGATG ACCCAAATC CACTCTCCCT	600
GCCTGTCAGT CTTGGAGATC AAGCCTCCAT CTCTGCAGG GCTAGTCAGA GCCTTGATACA	660
CAGTAATGGA AACACCTATT TACATTGGTA CCTGCAGAAG CCAGGCCAGT CTCCAAAGCT	720
CCTGATCTAC AGAGTTTCCA ACCGATTTTC TGGGGTCCCA GACAGGTTCA GTGGCAGTGG	780
ATCAGGGACA GATTTACAC TCAAGATCAG CAGAGTGGAG GCTGAGGATC TGGGAGTTTA	840
TTTCTGCTCT CAAAGTACAC ATGTTCTCTC GACGTTCCGT GGAGGCACCA AGCTGGAAAT	900
CAAACGACG CCAGTGTGAG AAAACAGCT GCGGAGGTG GTCGCGAATA CGATTACCCC	960
GCTGATGAAA GCCCAGTCTG TTCCAGGCAT GCGGTTGGCC GTTATTTATC AGGGAAAACC	1020
GCACTATTAC ACATTTGGCA AGGCCGATAT CGCGCGCAAT AAACCCGTTA CGCCTCAGAC	1080
CCTGTTGAG CTGGGTCTA TAAGTAAAC CTTACCCGGC GTTTTAGGTG GGGATGCCAT	1140
TGCTCGCGGT GAAATTTGCG TGGACGATGC GGTGACCAGA TACTGGCCAC AGCTGACGGG	1200
CAAGCAGTGG CAGGGTATTC GTATGCTGGA TCTCGCCACC TACACCGCTG GCGGCCTGCC	1260
GCTACAGGTA CCGGATGAG TCACGGATAA CGCCTCCCTG CTGCGCTTTT ATCAAAACTG	1320
GCAGCCGAG TGGAAGCCTG GCACAACGCG TCTTTACGCC AACGCCAGCA TCGGTCTTTT	1380
TGGTGCGCTG GCGGTCAAAC CTTCTGGCAT GCCCTATGAG CAGGCCATGA CGACGCGGGT	1440
CCTTAAGCCG CTCAAGCTGG ACCATACCTG GATTAACTG CCGAAAGCGG AAGAGGCGCA	1500
TTACGCCTGG GGCTATCGTG ACGGTAAAGC GGTGCGCGTT TCGCCGGGTA TGCTGGATGC	1560
ACAAGCCTAT GCGGTGAAAA CCAACGTGCA GGATATGGCG AACTGGGTCA TGGCAAACAT	1620
GGCGCCGAG AACGTTGCTG ATGCCTCACT TAAGCAGGGC ATCGCGCTGG CGCAGTCGCG	1680
CTACTGGCGT ATCGGGTCAA TGTATCAGGG TCTGGGCTGG GAGATGCTCA ACTGGCCCGT	1740
GGAGGCCAAC ACGGTGGTCG AGACGAGTTT TGGTAATGTA GCACTGGCGC CGTTGCCCGT	1800
GGCAGAAGTG AATCCACCGG CTCCCCCGGT CAAAGCGTCC TGGGTCCATA AAACGGGCTC	1860
TACTGGCGGG TTTGGCAGCT ACGTGGCCTT TATTCCTGAA AAGCAGATCG GTATTGTGAT	1920

GCTCGCGAAT ACAAGCTATC CGAACCCGGC ACGCGTTGAG GCGGCATACC ATATCCTCGA 1980

GGCGCTACAG TAGACTAGTG AATTCGAGCT 2010

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 442 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Met	Lys	Tyr	Thr	Ala	Ala	Ala	Gly	Ala	Ala	Ala	Met	Ala	Val	Ser	Gly	1	5	10	15
Ser	Val	Lys	Ser	Thr	Ser	Thr	Cys	Ser	Val	Thr	Gly	Asp	Ser	Thr	Ser	20	25	30	
Gly	Tyr	Trp	Asn	Trp	Arg	Lys	Gly	Asn	Lys	Tyr	Met	Gly	Tyr	Ser	Asp	35	40	45	
Ser	Gly	Thr	Tyr	Tyr	Asn	Ser	Lys	Ser	Arg	Ser	Thr	Arg	Asp	Thr	Ser	50	55	60	
Lys	Asn	Tyr	Tyr	Asn	Val	Thr	Ala	Asp	Thr	Ala	Thr	Tyr	Asn	Cys	Ala	65	70	75	80
Arg	Arg	Thr	Ala	Thr	Tyr	Tyr	Ala	Met	Asp	Tyr	Trp	Gly	Gly	Thr	Ser	85	90	95	
Val	Thr	Val	Ser	Ser	Gly	Ser	Thr	Ser	Gly	Ser	Gly	Lys	Gly	Ser	Gly	100	105	110	
Gly	Ser	Thr	Lys	Gly	Asp	Val	Met	Thr	Thr	Ser	Val	Ser	Gly	Asp	Ala	115	120	125	
Ser	Ser	Cys	Arg	Ala	Ser	Ser	Val	His	Ser	Asn	Gly	Asn	Thr	Tyr	His	130	135	140	
Trp	Tyr	Lys	Gly	Ser	Lys	Tyr	Arg	Val	Ser	Asn	Arg	Ser	Gly	Val	Asp	145	150	155	160
Arg	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Thr	Lys	Ser	Arg	Val	Ala	Asp	165	170	175	
Gly	Val	Tyr	Cys	Ser	Ser	Thr	His	Val	Thr	Gly	Gly	Gly	Thr	Lys	Lys	180	185	190	
Arg	Thr	Val	Ser	Lys	Ala	Val	Val	Ala	Asn	Thr	Thr	Met	Lys	Ala	Ser	195	200	205	
Val	Gly	Met	Ala	Val	Ala	Val	Tyr	Gly	Lys	His	Tyr	Tyr	Thr	Gly	Lys	210	215	220	
Ala	Asp	Ala	Ala	Asn	Lys	Val	Thr	Thr	Gly	Ser	Ser	Lys	Thr	Thr	Gly	225	230	235	240
Val	Gly	Gly	Asp	Ala	Ala	Arg	Gly	Ser	Asp	Asp	Ala	Val	Thr	Arg	Tyr	245	250	255	
Trp	Thr	Gly	Lys	Trp	Gly	Arg	Met	Asp	Ala	Thr	Tyr	Thr	Ala	Gly	Gly	260	265	270	
Val	Asp	Val	Thr	Asp	Asn	Ala	Ser	Arg	Tyr	Asn	Trp	Trp	Lys	Gly	Thr				

34

275	280	285
Thr Arg Tyr Ala Asn Ala Ser Gly Gly Ala Ala Val Lys Ser Gly Met 290 295 300		
Tyr Ala Met Thr Thr Arg Val Lys Lys Asp His Thr Trp Asn Val Lys 305 310 315 320		
Ala Ala His Tyr Ala Trp Gly Tyr Arg Asp Gly Lys Ala Val Arg Val 325 330 335		
Ser Gly Met Asp Ala Ala Tyr Gly Val Lys Thr Asn Val Asp Met Ala 340 345 350		
Asn Trp Val Met Ala Asn Met Ala Asn Val Ala Asp Ala Ser Lys Gly 355 360 365		
Ala Ala Ser Arg Tyr Trp Arg Gly Ser Met Tyr Gly Gly Trp Met Asn 370 375 380		
Trp Val Ala Asn Thr Val Val Thr Ser Gly Asn Val Ala Ala Val Ala 385 390 395 400		
Val Asn Ala Val Lys Ala Ser Trp Val His Lys Thr Gly Ser Thr Gly 405 410 415		
Gly Gly Ser Tyr Val Ala Lys Gly Val Met Ala Asn Thr Ser Tyr Asn 420 425 430		
Ala Arg Val Ala Ala Tyr His Ala Thr Ser 435 440		

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 7 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Glu Val Leu Gln Leu Glu Ser
 1 5

WHAT IS CLAIMED IS:

1. A fusion polypeptide comprising antibody variable light and heavy region amino acid sequences specific for a melanoma associated antigen operatively linked to a β -lactamase.
2. The fusion polypeptide of claim 1, wherein the fusion polypeptide is operatively linked to a secretory leader sequence.
3. The fusion polypeptide of claim 2, wherein the secretory leader sequence is pel B.
4. The fusion polypeptide of claim 1, wherein the β -lactamase is *E. cloacae* P99 having one or more amino acid substitutions at amino acid residues 537-541.
5. The fusion polypeptide of claim 4, wherein the substitution is Thr Ser Phen Gly Asn (SEQ ID. NO: 2) .
6. The fusion polypeptide of claim 1, wherein the variable light and heavy region amino acid sequences are separated by a spacer.
7. The fusion polypeptide of claim 6, wherein the spacer is Gly Ser Thr Ser Gly Ser Gly Lys Pro Gly Ser Gly Glu Gly Ser Thr Lys Gly (SEQ ID. NO: 13).
8. The fusion polypeptide of claim 1, wherein the polypeptide specifically binds a tumor cell antigen.
9. The fusion polypeptide of claim 8, wherein the tumor cell antigen is p97.
10. The fusion polypeptide of claim 1, wherein the fusion polypeptide has the amino acid sequence as shown in Figs. 4A and 4B (SEQ ID. NO: 20).

11. An isolated nucleic acid encoding the fusion polypeptide of claim 1.

12. A method for the delivery of a cytotoxic agent to a tumor cell comprising the administration of a fusion polypeptide comprising antibody variable light and heavy region sequences operatively linked to a β -lactamase, wherein the fusion polypeptide is reactive with a tumor cell antigen and the fusion polypeptide converts a prodrug to a cytotoxic drug, and the administration of an effective amount of the prodrug.

13. The method of claim 12, wherein the prodrug is derived from phenyleneamine mustard, doxorubicin, mitomycin C, paclitaxel, a vinca alkaloid, or melphalan.

14. The method of claim 12, wherein the tumor cell antigen is p97.



15. The method of claim 12, wherein the β -lactamase is *E. cloacae* P99 having one or more amino acid substitutions at residues 537-541.

16. The method of claim 15, wherein the amino acid substitutions are Thr Ser Phen Gly Asn (SEQ ID. NO: 2).

17. The method of claim 12, wherein the antibody variable light and heavy region sequences are separated by a spacer.

18. The method of claim 17, wherein the spacer is Gly Ser Thr Ser Gly Ser Gly Lys Pro Gly Ser Gly Glu Gly Ser Thr Lys Gly (SEQ ID. NO: 13).

1/11



2/11

FIG. 2




FIG. 7




FIG. 4A

[illegible]

4/11

FIG. 4B

CCITGTCGAGCTGGGTCTATAAGTAAACCTTACCCGGCGTTTTAGGTGGGATGCCATTGCTCGCGGTGAATTTTCGCTGGACGATGC
 309 L F E L G S I S K T F T G V L G G D A I A R G E I S L D D A
 GGTGACCAGATACITGGCCACAGCTGACGGGCAAGCAGTGGCAGGGTATTGCTATGCTGGATCTGCCACCTACACCGCTGGCGGCTGCG
 339 V T R Y W P Q L T G K Q W Q G I R M L D L A T Y T A G G L P
 GCTACAGGTACCGGATGAGGTCACGGATAACGCCCTCCCTGCTGGCTTTTATCAAACTGGTAGCCGCGAGTGGAAAGCCITGGCACAAACGCG
 369 L Q V P D E V T D N A S L L R F Y Q N W Q P Q W K P G T T R
 TCTTACGCCAACGCCAGCATCGGTCTTTTGGTGGCTGGCGGTCAAACTTCTGGCATGCCCTATGAGCAGGCGCATGACGACGCGCGGT
 399 L Y A N A S I G L F G A L A V K P S G M P Y E Q A M T T R V
 CCTTAAGCCGCTCAAGCTGGACCATACCTGGATTAACTGCCGAAAGCGGAAGAGCGGCATTACGCCITGGGGCTATCGTGACGGTAAAGC
 429 L K P L K L D H T W I N V P K A E A H Y A W G Y R D G K A
 GGTGCGGTTTCGCCGGTATGCTGGATGCACAAGCCTATGGCGTGAAACCAACGTCAGGATATGCCGAACCTGGTCAATGGCAAAACAT
 459 V R V S P G M L D A Q A Y G V K T N V Q D M A N W V M A N M
 GCGCGCGGAGAACGTTGCTGATGCCCTCACCCTAACGAGCGGCAATCGCGCTAGTGGCGTACTGGCGTATCGGGTCAATGATCAGGG
 489 A P E N V A D A S L K Q G I A L A Q S R Y W R I G S M Y Q G
 TCTGGGCTGGAGATGCTCAACTGGCCGCTGGAGGCCAACACGGTGGTCTGAGGcgaggttttgtaatGTAGCACTGGCGCGCTTGGCCGT
 519 L G W E M L N W P V E A N T V V E T S F G N V A L A P L P V
 GGCAGAGTGAATCCACCGGCTCCCCCGGTCAAAGCGTCCCTGGGTCCATAAAACGGGCTCTACTGGCGGTTTGGCAGCTACGTGGCCCTT
 549 A E V N P P A P P V K A S W V H K T G S T G G F G S Y V A F
 TATTCCTGAAAGCAGATCGGTATTGTGATGCTCGGAATACAAGCTATCCGAACCGGCACGCGTTGAGCGGCATACCATATCctcgo
 579 I P E K Q I G I V M L A N T S Y P N P A R V E A A Y H I L E
 ggcgctacagtagactagtgaaTTCGAGCT
 609 A L Q * T S E F E

5/11

FIG. 5A





FIG. 5B



6/11

FIG. 6A




FIG. 6B






FIG. 6C



7/11

FIG. 6D



8/11

FIG. 8







FIG. 9A



9/11

**FIG. 10A**

10/11

FIG. 10B






FIG. 10C



11/11

FIG. 10D



INTERNATIONAL SEARCH REPORT

International application No.
PCT:US98:08840

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07K 16 00, 17 00; C07H 21:02, 21:04; A61K 39:395

US CL : 530-391.1, 391.5, 391.7; 536-23.5; 424-130.1, 133.1, 138.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530:391.1, 391.5, 391.7; 536:23.5; 424:130.1, 133.1, 138.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CAS ONLINE

search terms: lactamase, antibod?, fusion

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	RODRIGUES, M.L. et al. Development of Humanized Disulfide-stabilized Anti-p185 ^{HER2} Fv-beta-Lactamase Fusion Protein for Activation of a Cephalosporin Doxorubicin Prodrug. Cancer Research. 01 January 1995, Vol. 55, pages 63-70, see abstract.	1-2, 6, 8, 12, 13, 17
X	KERR, D.E. et al. Regressions and Cures of Melanoma Xenografts following Treatment with Monoclonal antibody beta-Lactamase Conjugates in combination with Anticancer Prodrugs. Cancer Research. 15 August 1995, Vol. 55, pages 3558-3563, see abstract.	1-2, 6, 8-9, 12-14, 17

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claims; or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

13 JULY 1998

Date of mailing of the international search report

02 SEP 1998

Name and mailing address of the ISA US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

SHEELA J. HUFF

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/08840

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GOSHORN, S.C. et al. Genetic Construction, Expression, and Characterization of a Single chain Anti-Carcinoma Antibody Fused to beta-Lactamase. Cancer Research. 01 May 1993, Vol. 53, pages 2123-2127, see pages 2123-2124.	1-3, 6, 8, 11
X ----- Y	DE SUTTER, K. et al. A Bifunctional Murine:Human chimeric Antibody with one Antigen-Binding arm replaced by bacterial beta-Lactamase. Molecular Immunology, 1994, Vol. 31, pages 261-267, see abstract.	1, 11 ----- 12-13
X,P	SIEMERS, N.O. et al. Construction, Expression and Activities of L49-sFv-Beta-lactamase, a single-Chain Antibody Fusion Protein for Anticancer Prodrug Activation. Bioconjugate Chemistry. 1997. Vol. 8, pages 510-519, see abstract.	1-4, 6, 8-9, 11-12, 14, 17